

MAIN PANEL

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bayer Nematicur
Granular Nematicide

ACTIVE CONSTITUENT: 50 g/kg FENAMIPHOS
(an anticholinesterase compound)

For control of certain nematodes in tomatoes, crucifers and
ornamentals
in the home garden

(Yates bar)

300 g Net

Ancillary panel

How to use

Plants	Pests	Rate	How to apply
Tomatoes Herbaceous ornamentals (eg gerbera, dahlia, carnation, gladioli, chrysanthemum)	Root-knot nematode (not Tas.)	25 g/m ²	Apply within 7 days before seeding or transplanting. Sprinkle granules evenly over soil, lightly rake in, then water.
Crucifers (eg cabbage, cauliflower, broccoli, turnip, swedes)	Sugar beet nematode (not Tas.)		
Woody ornamentals (eg roses, camellia, hibiscus, ornamental peach)	Root-knot nematode (not Qld, Tas.)	25-50 g/m ²	Apply before planting and again each spring and autumn. Sprinkle evenly, lightly rake, then water. Use higher rate for heavy infestations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED

Nemacur Granular Nematicide offers control of nematodes in root systems, by the application of granules to the soil instead of spraying or dusting.

(diagram)

When applied at 25 g/m², Nemacur Granular Nematicide looks like this on the soil surface.

Caution

DO NOT allow chemical or container to get into drains, sewers, streams or ponds.

Storage and Disposal

Store in the closed, original container in a cool, dry place out of reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

Safety Directions

Very dangerous. Product is poisonous if absorbed by skin contact, inhaled or swallowed. Avoid contact with eyes and skin. Do not inhale dust. Wash hands after use.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre (131126). If swallowed, give one atropine tablet every 5 minutes until dryness of the mouth occurs. If poisoned by skin contact or through lungs, remove contaminated clothing and wash skin thoroughly and give atropine tablets as above. Get to a doctor or hospital quickly.

Ancillary panel (contd)

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

NemaCur® is a registered trademark of Bayer

NRA Approval Number 33291/0103

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
391-393 Tooronga Road
Hawthorn East Victoria 3123

Marketed and recommended by Arthur Yates & Co Limited
21a Richmond Road Homebush NSW 2140
www.yates.com.au

Batch Number:

Date of Manufacture: