

## READ SAFETY DIRECTIONS

**Confidor® Garden**

## INSECTICIDE

Active Constituent: 50 g/kg IMIDACLOPRID

For the systemic control of pests on flowers, shrubs, trees, fruit trees and vegetables in the home garden

**HOW TO PREPARE****Mixing**

Add the required number of sachets of Confidor to water in the sprayer and mix thoroughly. Let stand for 1 minute and then stir again. The prepared spray must be used on the day it is mixed. (1 sachet contains 5 g of Confidor).

5 L of spray mixture should be sufficient to thoroughly spray 2 fruit trees up to 3 m high, or to spray 30 large tomato plants.

**Caution**

DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

DO NOT spray if bees are feeding on flowering plants. Will kill bees.

**STORAGE AND DISPOSAL**

Keep out of the reach of children. Store in the closed, original container in a cool, dry place. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

**SAFETY DIRECTIONS**

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

**LIABILITY**

This product must be used strictly as directed. Bayer CropScience Pty Ltd may not be liable for loss or damage arising from failure to follow directions for use.

Confidor® is a registered trademark of Bayer

NRA Approval Number 49395/0201

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

**HOW TO USE**

Do not spray if it is raining or about to rain.

Do NOT spray when soil is dry and plants are suffering from moisture stress.

For best results spray in the morning when there is little or no wind and little direct heat from the sun.

PLANT	PEST	RATE	HOW TO APPLY
Roses	Aphids	Mix 1 sachet in 5 litres of water	Spray thoroughly at first sign of pests. Ensure underside of leaves are sprayed. For psyllids, spray at first sign then a week later.
Shrubs, plants and ornamental trees, including non-bearing and ornamental citrus	Aphids		
	Azalea lace bug		
	Citrus mealybug		
	Greenhouse thrips		
	Psyllids	Mix 2 sachets in 5 litres of water	Spray buds and flowers as needed.
	Sap-sucking bugs (including bronze orange bug)		
	Hibiscus flower beetle		
	Longtailed mealybug	Mix 2 sachets in 5 litres of water	Apply 3 sprays 2 weeks apart. Add a wetting agent.
	Soft scales	Mix 1 sachet in 5 litres of water	Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Peach, plum, nectarine, apricot and other stone fruit; capsicum; zucchinis, melons and other cucurbits; egg plant; potatoes; tomatoes	Green peach aphid	Mix 1 sachet in 5 litres of water	Spray thoroughly at first sign of pests. Ensure underside of leaves are sprayed.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED**

**DO NOT PICK STONE FRUIT FOR 21 DAYS AFTER SPRAYING**

**DO NOT PICK CAPSICUM, POTATOES OR EGG PLANTS FOR 7 DAYS AFTER SPRAYING**

**DO NOT PICK TOMATOES FOR 3 DAYS AFTER SPRAYING**

**DO NOT PICK CUCURBITS FOR 1 DAY AFTER SPRAYING**