

READ SAFETY DIRECTIONS

**Confidor® Plant**

INSECTICIDE

Active Constituent: 0.25 g/kg IMIDACLOPRID

For the systemic control of insect pests on flowers, shrubs, trees, fruit trees and vegetables in the home garden

**PRECAUTION**

Keep out of the reach of children.  
 DO NOT spray directly on humans, pets, exposed food, food utensils or food preparation surfaces.  
 DO NOT spray when bees are feeding on flowering plants.  
 FLAMMABLE. DO NOT store or use near fire or flame.

**BEWARE:** Deliberately sniffing or inhaling concentrated spray can be harmful or fatal.

**SAFETY DIRECTIONS**

May irritate the eyes and skin. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. Wash hands after use.

**STORAGE AND DISPOSAL**

Keep in a cool place out of the sun. DO NOT puncture or incinerate can, even when empty. Recycle empty cans if a facility is available or place used can in household rubbish.

**LIABILITY**

This product must be used strictly as directed. Bayer CropScience Pty Ltd may not be liable for loss or damage arising from failure to follow directions for use.

Confidor® is a registered trademark of Bayer

NRA Approval Number 49394/0201



FOR 24 HOUR SPECIALIST ADVICE  
 IN EMERGENCY ONLY  
 PHONE 1800 033 111

**HOW TO USE**

Do NOT spray if it is raining or about to rain.  
 Do NOT spray when soil is dry and plants are suffering from moisture stress.

PLANT	PEST	HOW TO APPLY
Roses	Aphids	Spray foliage thoroughly from a distance of 30 cm when insects are first seen or if insects re-appear. Spray underneath the leaves to kill pests which may be hiding out of sight. For hibiscus flower beetle, spray buds and flowers as needed. For longtailed mealybug, apply 3 sprays 2 weeks apart. For psyllids, spray at first sign and then a week later. For soft scales, spray in late spring or when small scales are first seen. Apply 3 sprays, 2 weeks apart. For best results spray in the morning when there is little or no wind and little direct heat from the sun. Avoid spraying new growth in full sun. For best results, move indoor potted plants to an outside shaded area, spray thoroughly and allow to dry. Slight spotting of sensitive ferns may occur. Avoid spraying any other surfaces as marking may occur.
Shrubs, flowers and ornamental trees, including non-bearing and ornamental citrus	Aphids Azalea lace bug Greenhouse thrips Hibiscus flower beetle Mealybugs Psyllids Sap-sucking bugs (including bronze orange bug) Soft scales	
Peach, plum, nectarine, apricot and other stone fruit; capsicum; zucchinis, melons and other cucurbits; egg plant; potatoes; tomatoes	Green peach aphid	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED**

**DO NOT PICK STONE FRUIT FOR 21 DAYS AFTER SPRAYING**  
**DO NOT PICK CAPSICUM, POTATOES OR EGG PLANTS FOR 7 DAYS AFTER SPRAYING**  
**DO NOT PICK TOMATOES FOR 3 DAYS AFTER SPRAYING**  
**DO NOT PICK CUCURBITS FOR 1 DAY AFTER SPRAYING**