

READ SAFETY DIRECTIONS

Confidor[®] Ready to Use

INSECTICIDE

Active Constituent: 0.125 g/L IMIDACLOPRID

For the systemic control of insect pests on flowers, shrubs, trees, fruit trees and vegetables in the home garden

HOW TO PREPARE

The product is ready to use. Shake well before use.

CAUTION

DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.
DO NOT spray if bees are feeding on flowering plants. Will kill bees.

STORAGE AND DISPOSAL

Keep out of the reach of children. Store in the closed, original container in a cool, dry place. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Wash hands after use.

LIABILITY

This product must be used strictly as directed. Bayer CropScience Pty Ltd may not be liable for loss or damage arising from failure to follow directions for use.

Confidor[®] is a registered trademark of Bayer

NRA Approval Number 50547/0201

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

HOW TO USE

Do NOT spray if it is raining or about to rain.

Do NOT spray when soil is dry and plants are suffering from moisture stress.

PLANT	PEST	HOW TO APPLY
Roses	Aphids	Spray foliage thoroughly when insects are first seen or if insects re-appear. Spray underneath the leaves to kill pests which may be hiding out of sight. For hibiscus flower beetle, spray buds and flowers as needed. For longtailed mealybug, apply 3 sprays 2 weeks apart. For psyllids, spray at first sign and then a week later. For soft scales, spray in late spring or when small scales are first seen. Apply 3 sprays, 2 weeks apart. For best results spray in the morning when there is little or no wind and little direct heat from the sun. Avoid spraying blooms or flowers in full sun as slight petal marking may occur.
Shrubs, plants and ornamental trees, including non-bearing and ornamental citrus	Aphids Azalea lace bug Greenhouse thrips Hibiscus flower beetle Mealybugs Psyllids Sap-sucking bugs (including bronze orange bug) Soft scales	
Peach, plum, nectarine, apricot and other stone fruit; capsicum; zucchinis, melons and other cucurbits; egg plant; potatoes; tomatoes	Green peach aphid	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED

DO NOT PICK STONE FRUIT FOR 21 DAYS AFTER SPRAYING

DO NOT PICK CAPSICUM, POTATOES OR EGG PLANTS FOR 7 DAYS AFTER SPRAYING

DO NOT PICK TOMATOES FOR 3 DAYS AFTER SPRAYING

DO NOT PICK CUCURBITS FOR 1 DAY AFTER SPRAYING