

MAIN PANEL

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Gauntlet[®]

INSECTICIDE

ACTIVE CONSTITUENT: 25 g/L BETACYFLUTHRIN

GROUP 3A INSECTICIDE

2.5 L

For the control of various turf insect pests and for control of certain nuisance pests around and inside buildings as specified in the directions for use.

IMPORTANT: Read the attached booklet before use.

Front page of booklet

CAUTION
KEEP OUT REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GAUNTLET INSECTICIDE
ACTIVE CONSTITUENT: 25 G/L BETACYFLUTHRIN

GROUP	3A	INSECTICIDE
--------------	-----------	--------------------

For the control of various turf insect pests and for control of certain nuisance pests around and inside buildings as specified in the directions for use.

STORAGE AND DISPOSAL Store in tightly closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.
The method of disposal depends on the container type, refer to the label attached to the container for full disposal directions.

SAFETY DIRECTIONS

When using the prepared spray by hand-held equipment, wear cotton overalls buttoned to the neck and wrist (worn over normal clothing) and a washable hat and elbow length PVC gloves. After each days use wash gloves and contaminated clothing.

RE-ENTRY

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist and chemical resistant gloves. Clothing must be laundered after each days use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 131126) Remove from contaminated area. Apply artificial respiration if not breathing.

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

IMPORTANT: READ THIS BOOKLET BEFORE USE

APVMA Approval Number: 59425/2.5/0905

Gauntlet® is a registered trademark of Bayer.

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800
Website: www.bayercropscience.com.au

Batch Number:

DOM:

BOOKLET**Directions for use:**

Situation	Rate	Pest	Critical Comments
Turf	20 - 40 mL/ 100 m ² or 2 - 4 L/ha or 250 - 500 mL/bowling green	Armyworm Cutworm Webworm	Apply to mown turf in a minimum application volume of 200 L/ha (or 2 L/100 m ²) and irrigate treated areas after application with up to 4 mm water. Use the higher rate when pest numbers are very high. Repeat as necessary. Application after mowing is recommended to minimise loss of insecticide in clippings.
	40 – 60 mL/100 m ² or 4 – 6 L/ha or 500 – 750 mL/bowling green	Adult Billbug, Adult Argentine Stem Weevil, Adult Scarabs, (including African black beetle), Ants	
Native plants	6 mL/L	Sawfly larvae	Apply thoroughly as a full cover spray. Repeat at beginning of renewed pest activity. See note on native plants. Do not spray if it is raining or about to rain.
Around and inside buildings and structures	10 mL/L	Ants, spiders, cockroaches, fleas, crickets	Apply as a coarse spray to infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Mixing: Add the required quantity of Gauntlet Insecticide to water while stirring.

Application: Do not spray the moving parts of any machinery, electric motors or switchgear with the water-based product.

Compatibility: Physical compatibility with other products has not been tested with this formulation. It is recommended that any mixtures should be tested prior to mixing commercial quantities.

PRECAUTIONS: For surface treatment only – do not spray into the atmosphere using foggers or other devices designed to create an aerosol.

Do not spray directly on humans, exposed food or food utensils or food preparation areas.

NOTE ON NATIVE PLANTS

Gauntlet Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole target group.

INSECTICIDE RESISTANCE WARNING

GROUP	3A	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management Gauntlet is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Gauntlet and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Gauntlet or other Group 3A insecticides are used repeatedly. The effectiveness of Gauntlet on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from failure of Gauntlet to control resistant insects.

For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science representative.

Protection of Livestock

DO NOT graze treated turf or feed grass clippings to livestock. Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

Protection of Wildlife, Fish, Crustacea and Environment: Dangerous to fish. Cover fish tanks before using. Do not contaminate ponds, waterways or drains with this product or used container.

Rear Panel (attached to container)

STORAGE AND DISPOSAL: Store in tightly closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

When using the prepared spray by hand-held equipment, wear cotton overalls buttoned to the neck and wrist (worn over normal clothing) and a washable hat and elbow length PVC gloves. After each days use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 131126) Remove from contaminated area. Apply artificial respiration if not breathing.

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet.

APVMA Approval Number: 59425/2.5/0905

Gauntlet® is a registered trademark of Bayer.

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800
Website: www.bayercropscience.com.au

Batch Number:

DOM:

